

UNITED STATES PATENT APPLICATION**COMPOSITIONS AND METHODS FOR THE
REPAIR AND CONSTRUCTION OF BONE AND
OTHER TISSUE****Inventor:** Chia-Nang (Sophia) Chang, M.D., Ph.D.**Filed by Hand at the United States Patent and Trademark Office on April 19, 2001**

1 **COMPOSITIONS AND METHODS FOR THE REPAIR AND**
2 **CONSTRUCTION OF BONE AND OTHER TISSUE**

3
4 **FIELD OF THE INVENTION**

5 The present invention relates to compositions and methods for repair and construction of
6 bone, cartilage, muscle, adipose and fibrous tissues, and the enhancement of healing of such
7 tissues.

8 **DESCRIPTION OF THE RELATED ART**

9 Over one million procedures in the United States each year involve bone and cartilage
10 replacement (Langer, R., et al., 1993, *Tissue Engineering*, SCIENCE. 920:260-266.1). The
11 current supply of bone marrow does not meet the demand for bone grafts. Sequences of various
12 bone morphogenic proteins (BMP) are known and have been successfully cloned. Since the
13 genes of the BMP family were successfully cloned, large quantities of individual BMP proteins
14 have been made available by recombinant-DNA technology in hope of assisting bone formation.
15 For example, Yasko et al. first demonstrated the effects of recombinant human BMP-2 on bone
16 formation in the rat segmental femoral-defect model (Yasko, A.W., et al., *The Healing of*
17 *Segmented Defects, Induced by Recombinant Human Bone Morphogenetic Protein (rhBMP-2): a*
18 *Radiographic, Histological, and Biomechanical Study in Rats*, J. BONE JOINT SURG. 74:659-670
19 (1992)). Studies of this nature have been performed in the sheep femur (Gerhart, T.N., et al.,
20 *Healing Segmented Femoral Defects in Sheep Using Recombinant Human Bone Morphogenetic*
21 *Protein*, CLIN. ORTHOP. 293:317-326 (1993)), and in canine spine and mandible models.
22 Although the results are encouraging in studies in which BMP protein is implanted directly to a
23 defective area, many disadvantages continue to exist. For example, a large amount of protein (up
24 to milligram quantities) is often required to stimulate significant new bone formation *in vivo*.
25 This requirement for large quantities of the protein increases the risk of unwanted side effects.

Moreover, the lack of a system to deliver proteins in a continuous manner over time may further hamper this technique. (Cook, S.D., et al., *In vivo Evaluation of Recombinant Human Osteogenic Protein Implants as a Bone Graft Substitute for Spinal Fusions*, SPINE 19:1655-1663 (1994).

Additionally, autologous or autogenous vascularized bone grafts may be harvested from temporal, scapular, radial, rib, iliac, fibular, metatarsal bones, but they have the following shortcomings: (1) Grafts are limited in supply; (2) the bone regenerated from peptides, (rhBMPs) demineralized bone powder, or combinations of both are small in size (Zegzula, H.D., et al., 1997, *Bone Formation with Use of rhBMP-2 (Recombinant Human Bone Morphogenetic Protein (rhBMP-2))*, J. BONE JOINT SURG. AM. 79(12):1778-1790; Nicderwanger, M., et al., 1996, *Demineralized Bone Matrix Supplied by Bone Banks for a Carrier of Recombinant Human Bone Morphogenetic Protein (rhBMP-2: A Substitute for Autogenic Bone Grafts* J. ORAL IMPLANTOL, 22(3-4):210-215; and (3) allografts are limited in usage because of immunological rejections, transmission of infectious diseases, premature resorption, and donor shortage. (Zegzula, H.D., et al., 1997; Rabies A.B., Chay, S.H., et al., 2000, *Healing of Autogenous Intramembranous Bone in the Presence and Absence of Homologous Demineralized Intramembranous Bone*, AM. J. ORTHOD. DENTOFACIAL. ORTHO. p. 117(3):288-297; Niederwanger, M., et al., 1996; Furukawa, T., et al., 2000, *Histomorphometric Study on High-Strength Hydroxyapatite/poly (L-lactide) Composite Rods for Internal Fixation of Bone Fractures*, J. BIOMED. MATER. RES. 5;50(3):410-419). Consequently, although BMP-2 has been known for 35 years, it is still difficult to use BMP to repair large size defects due to these and other disadvantages.

One approach to repair large defects is to combine polymers with BMP. In recent years, tissue engineering has had some success in employing biocompatible, biodegradable polymers

1 seeded with living cells to create functional tissue. (Langer,R., et al., 1993, *Tissue Engineering*,
2 SCIENCE. 920; 260-266). The field of tissue engineering has been developing to overcome the
3 problem of donor-tissue scarcity of either whole organs or reconstructive grafts. Earlier work
4 with osteoblasts and PGA/PLA fiber meshes showed that osteoblasts-PGA/PLA constructs can
5 be fabricated and, further, that when they are implanted into an animal host, new bone
6 composition is generated with the final morphology established by the shape of the polymer
7 scaffold.

8 It is believed that osteoblasts are derived from pluripotential mesenchymal stem cells and
9 may develop into a lineage capable of producing bone, muscle, cartilage, adipose and fibrous
10 tissue (Metcalf, D. 1989, *The Molecular Control of Cell Division, Differentiation Commitment*
11 *and Maturation in Haemopoietic Cells*, NATURE 339:27-30); Wiles, M.V., et al., 1991; *Multiple*
12 *Hematopoietic Lineages Develop From Embryonic Stem (ES) Cells in Culture*, DEVELOPMENT
13 111:259-267). The ultimate fate of an undifferentiated stem cell is determined by local growth
14 factors and by the environment to which the mesenchymal stem cell is exposed (Goshima, J.,
15 Goldberg V.M., et al., 1991, *The Origin of Bone Formed in Composite Grafts of Porous Calcium*
16 *Phosphate Ceramic Loaded With Marrow Cells*, CLIN. ORTHOP. 269:274-283).

17 Bone morphogenic protein (BMP) has been shown to encourage the development of
18 osteoblasts from mesenchymal stem cells. (Lou, J., et al., 1999, *Gene Therapy: Adenovirus-*
19 *Mediated Human Bone Morphogenetic Protein-2 Gene Transfer Induces Mesenchymal*
20 *Progenitor Cell Proliferation and Differentiation In Vitro and Bone Formation In Vitro*. J.
21 ORTHO. RES. 17:43-50).

22 It was previously demonstrated that BMP-2 gene transfer into mesenchymal progenitor
23 cells could induce undifferentiated stem cells *in vitro* to follow an osseous lineage resulting in

1 bone formation, although continued incubation for up to 30 days and several medium changes
2 were required. (Lou, J., et al., 1999, *Gene Therapy: Adenovirus-Mediated Human Bone*
3 *Morphogenetic Protein-2 Gene Transfer Induces Mesenchymal Progenitor Cell Proliferation*
4 *and Differentiation In Vitro and Bone Formation In Vitro*, J. ORTHO. RES. 17:43-50). However,
5 it was heretofore unknown as to whether the application of such gene therapy *in vivo* as disclosed
6 and claimed herein would indeed be effective in repairing bone and other tissues.

7 SUMMARY OF THE INVENTION

8 The *in vivo* experiments disclosed herein demonstrate that bone and other tissue
9 repair is induced and enhanced by adenovirus BMP-2 transduction. One aspect of the invention
10 relates to novel compositions comprising genetically engineered cells and one or more polymers.
11 In an additional aspect, the present invention relates to a method for repairing tissue, for
12 example, cranioskeletal or maxillary bone defects, comprising transducing the BMP-2 gene into
13 bone marrow stromal cells which are harvested from a subject, combining the genetically
14 engineered cells with at least one polymer, and implanting the combination at the site of the
15 defect. The BMP-2 protein is advantageously produced as long as the transduced gene stays in
16 the cells.

17 BRIEF DESCRIPTION OF THE DRAWINGS

18 Fig. 1 is a schematic depicting a method of the present invention.

19 Fig. 2 is a series of photographs as follows. 2A (Left upper): The maxilla bone in minipigs was
20 exposed by removal of periosteum. 2B (Right upper): The maxilla bony defect in minipigs: 3 x
21 1.2 cm² bilaterally. 2C (Left lower): One L-shape resorbable plate was wired on the defect edge
22 to secure the space from soft tissue invagination. 2C (Right lower): The defect was filled by the
23 gene engineered bone marrow mesenchymal stromal cells.

- 1 Fig. 3 is a series of photographs of the maxilla after 3 months implantation of the composition
2 of the invention *in vivo*. Left: control group depicting lack of bone formation. Right:
3 experimental group depicting bone formation and bridging of the defect.
- 4 Fig. 4 is a 3D CT image of maxilla repair, 3 months *in vivo* (Left: control, Right: experiment).
- 5 Fig. 5 are photomicrographs of H&E and Von Kossa stainings. A. The H&E staining of the
6 control site showed no bony formation (40x). B. The H&E staining of the experimental site
7 showed cancellous bone formation (40x). C. The Von Kossa staining of the BMP-2 site showed
8 lamellar bone formation (40x). D. The higher power field of the BMP-2 shows bone formation.
- 9 Fig. 6 is a Western blot of BMP-2 expression in MSCs.
- 10 Fig. 7 is a series of photographs of the cranial defect in minipigs: $2 \times 5 \text{ cm}^2$. Left upper: the
11 bony defect bilaterally. Right upper: the underlying dura were removed. Lower: the Alloderm
12 was used for dura repair.
- 13 Fig. 8 is a series of photographs depicting the cranium in Group I after 3 months implantation *in*
14 *vivo*. A. View from cranium. B. Transverse view.
- 15 Fig. 9 is a photograph depicting the cranium in Group IV after 6 weeks (A). 3 months (B).
16 implantation *in vivo*. Transverse view 3 months (C).
- 17 Fig. 10 is a 3D CT image of Group I at 3 months (Left: BMP-2 site, Right: β Gal site).
- 18 Fig. 11 is a 3D CT images of Group II at 3 months (Left: BMP-2 site, Right: β Gal site).
- 19 Fig. 12 is a 3D CT image of Group III at 3 months (Left: BMP-2 site, Right: β Gal site).

Fig. 13 is a series of 3D CT images. 13A is a 3D CT image of Group IV at 6 weeks (Left: BMP-2 site, Right: β Gal site). 13B is a 3D CT image of Group IV at 3 months (Left: BMP-2 site, Right: β Gal site).

Fig. 14 is a series of photographs of the H & E staining. 14A is a H & E staining of the experimental site showed cancellous bone formation (200x). 14B is a H & E staining of the control site showed no bone formation (200x).

Fig. 15 is a series of photographs of the Von Kossa staining of the BMP-2 site. 15A depicts lamellar bone formation (50x). 15B depicts higher power field of the bone formation at the BMP-2 site.

Fig. 16 is a series of photographs of the immunochemical stain for detecting adenovirus in Group 4. 16A: positive in pre-implantation culture (200X). 16B: negative in 6 weeks; and 16C: 3 months samples (200X).

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

While the present invention may be embodied in many different forms, several specific embodiments are discussed herein with the understanding that the present disclosure is to be considered only as exemplification of the principles of the invention, and it is not intended to limit the invention to the embodiments illustrated.

It will be apparent to those skilled in the art that various modifications and variations can be made in the methods and compositions of the present invention without departing from the spirit or scope of the invention. Thus, it is intended that the present invention include modifications and variations that are within the scope of the appended claims and their equivalents.

1 The present invention is directed to methods, kits, combinations and pharmaceutical
2 compositions for treating or repairing bone and other tissue defects. Besides being useful for
3 human treatment, the present invention is also useful for veterinary treatment of mammals,
4 exotic animals and farm animals.

5 The use of the term "about" in the present disclosure means "approximately," and use of
6 the term "about" indicates that amounts and dosages slightly outside the cited ranges may also be
7 effective and safe, and such amounts and dosages are also encompassed by the scope of the
8 present claims.

9 The phrase "cell/polymer composition" or "pharmaceutical composition" or simply
10 "formulation" or "composition" refers to the combination of the genetically engineered cells
11 disclosed herein and at least one pharmaceutically acceptable polymer or carrier appropriate for
12 use as a pharmaceutical product.

13 The phrase "pharmaceutically acceptable" is used adjectivally herein to mean that the
14 modified noun is appropriate for use in a pharmaceutical product. Pharmaceutically acceptable
15 polymers and biomaterials to be used as a matrix or carrier for the therapeutic cells, and for
16 implanting into the defect include, but are not limited to, alginate (Sigma), alginate (Pronova),
17 alginate-RGD, and collagen I (Pancogen). Other pharmaceutically acceptable polymers and
18 biomaterials include any natural or synthetic polymers such as PDS, PLA, PGA, PALGA,
19 PLGA, hydroxyapatite, and the like.

20 The phrase "bone or other tissue defect" means a defect or injury in or to the bone,
21 cartilage, muscle, adipose or other fibrous tissue.

22 A "bone or other tissue defect-effective amount," or "effective amount" is intended to
23 qualify the amount of cell/polymer composition required to treat a bone or other tissue defect in a

1 mammal, or relieve to some extent one or more of the symptoms associated with, or related to, a
2 bone or other tissue defect in a mammal.

3 The compositions of the present invention are used in a "effective amount." This means
4 that the concentration of the BMP protein is such that a therapeutic level of protein is delivered
5 over the term that the cell/polymer composition is to be used. Such delivery is dependent on a
6 number of variables including the time period for which the individual dosage unit is to be used,
7 the flux rate of the therapeutic agent, for example, BMP protein, from the cells and polymer,
8 surface area of application site, etc. It is understood, however, that specific dose levels of the
9 therapeutic agents of the present invention for any particular patient depends upon a variety of
10 factors including the activity of the specific compound employed, the age, body weight, general
11 health, sex, and diet of the patient, the time of administration, the rate of excretion, the drug
12 combination, and the severity of the particular disorder being treated and form of administration.
13 Treatment dosages generally may be titrated to optimize safety and efficacy.

14 The phrase "combination therapy" embraces the administration of BMP producing cells
15 of various types, or employing the inventive formulations with other drugs or therapies. The
16 beneficial effect of the combination includes, but is not limited to, pharmacokinetic or
17 pharmacodynamic co-action resulting from the combination of therapeutic agents.
18 Administration of these therapeutic agents in combination typically is carried out over a defined
19 time period (usually minutes, hours, days, weeks, months or years depending upon the
20 combination selected). "Combination therapy" generally is not intended to encompass the
21 administration of two or more of these therapeutic agents as part of separate monotherapy
22 regimens that incidentally and arbitrarily result in the combinations of the present invention.
23 "Combination therapy" is intended to embrace administration of these therapeutic agents in a

1 sequential manner, that is, where each therapeutic agent is administered at a different time, as
2 well as administration of these therapeutic agents, or at least two of the therapeutic agents, in a
3 substantially simultaneous manner. Sequential or substantially simultaneous administration of
4 each therapeutic agent can be effected by any appropriate route including, but not limited to,
5 topical, oral routes, percutaneous routes, intravenous routes, intramuscular routes, and direct
6 absorption through mucous membrane tissues. The therapeutic agents can be administered by
7 the same route or by different routes. "Combination therapy" also can embrace the
8 administration of the therapeutic agents as described above in further combination with other
9 biologically active ingredients.

10 Examples of suitable pharmaceutically acceptable formulations are stated herein.
11 Additionally, drug formulations are discussed in, for example, Remington's The Science and
12 Practice of Pharmacy (2000) and Liberman, H.A. and Lachman, L., Eds., Pharmaceutical Dosage
13 Forms, Marcel Decker, New York, N.Y., 1980.

14 **EXAMPLE 1:**

15 In this example, the repair of critical size craniofacial bone defect in mini-pigs by BMP-2
16 gene engineered autologous bone marrow stromal cells (MSC) was assessed. The MSC were
17 separated from the aspirate from the iliac crest from 40 minipigs one month before implantation.

18 The general method is illustrated schematically in Fig. 1. As shown below, near-complete
19 critical-size craniofacial defect repair was achieved by simple aspiration of bone marrow stromal
20 cells and after one month implantation of tissue engineered constructs.

21 **MATERIALS AND METHODS**

22 **Construction of Recombinant Adenovirus**

23 Recombinant adenovirus BMP-2 was constructed by *in vivo* homologous recombination
24 in 293 cells between expression vector pAC-cytomegalovirus BMP-2 and the large Cl_a I

fragment (2.6-100 mm) of adenovirus type 5 genome DNA. Briefly, to make recombinant plasmid pAC-cytomegalovirus BMP-2, the human BMP-2 gene (No. 40345; ATCC, (American Type Culture Collection), Rockville, MD, U.S.A.) was inserted into the vector pAC cytomegalovirus. The resulting plasmid was made linear by Xho I and cotransfected with the large Cla I fragment of wild-type adenovirus type 5 genome DNA by lipofectamine (GIBCO BRL, Grand Island, NY, U.S.A.). For each 35-mm dish of 293 cells, 2 mg of each kind of DNA (total: 4 mg) was transfected. Plaques containing recombinant adenovirus were picked after 10 days of agarose overlay and incubation at 37°C. The viral stocks were propagated and titered in 293 cells. The recombinant adenovirus with human BMP-2 gene, which was under transcriptional control of the cytomegalovirus early gene promoter/enhancer, is known as adenovirus BMP-2 (Lou, J., et al., 1999, *Gene Therapy: Adenovirus-Mediated Human Bone Morphogenetic Protein-2 Gene Transfer Induces Mesenchymal Progenitor Cell Proliferation and Differentiation In Vitro and Bone Formation In Vitro*, J. ORTHO. RES. 17:43-50). The recombinant adenovirus containing the bacterial β -galactosidase (lac z) gene was constructed by the same procedure (Lou, J., et al. 1996, *Adenovirus-Mediated Gene Transfer into Tendon and Tendon Sheath*, J. ORTHOP. RES. 14:513-517), and the virus is known as adenovirus β -galactosidase. Viral stock of 293 lysate was titered in 293 cells by serial dilution and plaque-formation assay.

Western Blot Detection of Adenovirus BMP-2 Expression in MSC (Bone Marrow Stromal Cells) *In Vitro*

Cells from MSC were grown in Dulbecco's modified Eagle medium with 10% fetal calf serum to 90% confluence. The cells were infected with adenovirus β -galactosidase or adenovirus BMP-2 at a multiplicity of infection of 50. The infected cells were maintained in Dulbecco's modified Eagle medium with 10% fetal calf serum and 50 mg/ml ascorbic acid. The

1 medium was collected 7 days later and then immunoprecipitated with monoclonal antibody
2 h4b2/5.10.24 (a gift from Genetics Institute, Cambridge, MA, U.S.A.). The immunoprecipitation
3 was then run on a 12.5% sodium dodecyl phosphate-polyacrylamide gel electrophoresis and
4 transferred overnight to a polyvinylidene difluoride membrane (Millipore, Bedford, MA,
5 U.S.A.). The expressed protein was detected with monoclonal antibody h4b2/5.10.24 and goat
6 anti-mouse IgG alkaline-phosphatase conjugate (Sigma Chemical, St. Louis, MO, U.S.A.).

7 **Bone Marrow Aspiration and Adenoviral Infection**

8 40 minipigs (Mitsae Pig, CGU, Taiwan) were randomly assigned to four groups. The
9 four groups (n=10) of different polymers: Group I, Alginate (Pronova); Group II, Alginate
10 (Sigma: self-purified); Group III, Alginate (RGD) and Group IV Pancogene S (Gattefosse). The
11 minipigs were sacrificed at 6 weeks and at 3 months.

12 At the first stage of surgery, bone marrow aspiration was performed when minipigs were
13 under intramuscular injection of 1 ml/ 10 kg 2% Rompum (Bayer, USA), Ketamine injection
14 (50mg/ml, Yung Shin Pharmaceutical Industrial Co., Taiwan). 20 cc of the aspirated bone
15 marrow was mixed with 2 cc of heparin sodium (5000 unit/ml, Agglutex, China Chemical &
16 Pharmaceutical Co., Taiwan). Aspirated iliac crest marrow (5-10 ml) was transferred to sterile
17 tubes, to which 20 ml complete medium was added. The tubes were spun at 1000 rpm for 5 min
18 to pelletize the cells, the supernatants and fat layers were removed, and the cell pellets (2.5-5.0
19 ml) were resuspended and loaded onto 70% Percoll (Sigma, St. Louis, Missouri) gradients.
20 These gradients were centrifuged at 460 x g for 15 min and harvested into three fractions with a
21 pipet: the top 25% of the gradients (low density cells), pooled density = 1.03 g per ml; the
22 middle 50% of the gradients (high density cells), pooled density = 1.10 g per ml; density = 1.14 g
23 per ml. In preliminary experiments, each of these three pools was plated separately in complete
24 medium in 100 mm dishes. Adherent marrow-derived mesenchymal cells were observed to be

1 localized to the low-density fraction. To produce adherent cell cultures for all subsequent
2 experiments, only the cells in the low density fraction were plated. The medium was changed
3 every 3-4 days. After one month, an adequate number of cells was reached.

4 At the experimental site, adenovirus mediated human BMP-2 gene transfer to MSCs was
5 performed 5-7 days before the secondary stage of surgery. As a control, adenovirus mediated
6 β Gal gene transfer was carried out at the same time and for the same period of time. The cells
7 were trypsinized by 25% trypsin and washed for three times. Cell numbers were determined
8 with a hemocytometer. The cells were then mixed with four different polymers to achieve the
9 same concentration of $50 \times 10^6/\text{ml}$. After one hour in the incubator, the cells in group 4 with the
10 polymer Pancogene S became more gel-like (in the dishes). In groups 1, 2 and 3, the cells in
11 CaSO_4 /alginate were gels ten minutes after mixing.

12 **Open Implantation on the Cranial Defects After Creating Cranial Bony Defect**

13 At the secondary stage of surgery, all minipigs were intubated and on ventilators. Two
14 cranial defects of $2 \times 5 \text{ cm}^2$ each were created by removal of periosteum from above and dura
15 from below, ensuring that no osteoinductive tissue existed (Fig. 7). The dura defects were
16 replaced with acellular dermis (Pig Alloderm, LifeCell USA).

17 3ml of cell/polymer constructs were implanted to each site. The right sided defect was
18 covered with adenovirus-mediated BMP-2 gene transfered MSCs/polymer. The left side defect
19 was filled with adenovirus-mediated β Gal gene transfered on MSCs/polymer. The scalp wounds
20 were closed in a watertight manner.

21 **Histologic Examination of Bone - Samples Harvested from the Cranial Defects**

22
23 Harvested samples were fixed in buffered 10% formalin for 72 hours and then sawed in
24 half. Half of them were decalcified in Decalcifier I (Surgipath, Northbrook, IL, U.S.A.) solution
25 for another 48 hours. The samples then underwent procedures for embedding, incising, and

1 staining with hematoxylin and eosin. The nondecalcified other half of the samples was stained
2 with von Kossa's silver nitrate for demonstration of matrix mineralization (Heise, C., et al.,
3 1997).

4 **Immunohistochemistry for Adenovirus Hexon Protein**

5 Immunohistochemistry was performed using formalin-fixed, paraffin embedded tumor
6 sections. After removal from paraffin, slides were hydrated and digested with pronase (Meridian
7 Diagnostics, Cincinnati, OH). The primary antibody (MAB805, Chemicon International,
8 Temecule, CA) was applied at 1:500 dilution, and slides were incubated for 1 hr at 35°C. A
9 biotinylated goat anti-mouse secondary was then applied, followed by a streptavidin-horseradish
10 peroxidase conjugate. Diaminobenzidine (DAB) was used as the chromogen before slides were
11 counterstained with hematoxylin.

12 **Biomechanical Analysis**

13 The samples were stored at -70°C before testing. All samples of engineered bones were
14 sawed using a 6-mm dermal punch to form disks. Each disk was immediately mounted in an
15 electrically insulating cylindrical confining chamber. The chamber was mounted in a servo-
16 controlled Dynastat mechanical spectrometer (IMASS, Hingham, MA) interfaced to a computer
17 (Lou, J., et al., 1996, *Adenovirus-Mediated Gene Transfer into Tendon and Tendon Sheath*, J.
18 ORTHOP. RES. 14:513-517). The samples were equilibrated at room temperature in 0.15M PBS,
19 pH 7.4, containing 100 units/ml penicillin G, and 100 ug/ml streptomycin. Samples were
20 compressed between a porous polyethylene platen at the base of the chamber.

21 After mounting each disk in the confined compression chamber, the distance between the
22 porous platen and the chamber was decreased until a signal of ~5 gm was detected by the load
23 cell. This distance was taken to be the sample thickness. Each disk was compressed by 10
24 sequential increments of 2.5-3.0% static strain, up to a maximum of 25-30% total strain. After

1 each increment, the load was recorded every 0.5 seconds for 100 seconds. Stress relaxation data
2 was fit to a poroelastic model of material behavior that yielded values for the material properties;
3 equilibrium modulus.

4 **Statistical analysis**

5 All measurements were collected in triplicate and expressed as means \pm standard
6 deviations. Single factor analysis of variance (ANOVA) was employed to assess the statistical
7 significance of results for all the 3D CT images of the repaired bone areas.

8 **RESULTS**

9 **Recombinant Adenovirus Construction and Human BMP-2 Expression**

10 Adenovirus BMP-2, a replication-defective adenovirus vector encoding human BMP-2,
11 was constructed in applicants laboratory. In this vector, transcription of the BMP gene is
12 controlled by cytomegalovirus early gene promoter/enhancer. The recombinant adenovirus
13 encoding *Escherichia coli* lac z gene (adenovirus bgalactosidase) was constructed by the same
14 procedure as the control vector. These two recombinant adenoviruses have been deleted only in
15 the E1 region of the genome. The E3 region of the virus genome was purposefully retained intact
16 on the vectors because of evidence that the E3 proteins of the virus inhibit the host immune
17 response (Horwitz, M.S., et al., 1995, *Model System for Studying the Effects of Adenovirus E3*
18 *Gene on Virulence in Vivo*, CURR. TOPICS MICROBIOL. IMMUNOL. 199:195-211). Titered in 293
19 cells, the viral lysate of the vectors are usually 10^8 plaque-forming units per milliliter.

20 The gene expression of human BMP-2 protein in MSCs, which was mediated by
21 adenovirus BMP-2 gene transfer, was assessed by detecting the coding product with monoclonal
22 antibody h4b2/5.10.24 in transduced cells. The secreted active BMP-2 molecules were detected
23 in cultured medium from adenovirus BMP-2-transduced MSCs by immunoprecipitation and
24 immunoblot analysis but were not found in cultured medium from either adenovirus b-

1 galactosidase-infected MSCs. The BMP gene transfected by adenovirus seems well expressed
2 and processed in MSCs. The molecular weight of secreted BMP-2 indicated mature status
3 without prepeptide. The band was noticeable at 18KDa due to overlap of bands at 16 and 22.

4 Grossly, in group 1, 2, 3, there is no bone formation at cranial defects sites. In Group 1,
5 after 3 months of *in vivo* treatment, incomplete repair was detected (Fig. 8). There was
6 noticeable whitish bone formation at the site infected with BMP-2 from 6 weeks on (Fig. 9A)
7 while no obvious bone formation was found at the β Gal infection site (Fig. 9).

8 3D-CT image studies were performed for each sample. In group I, there were some
9 scattered bony formations at both reconstructed areas (Fig. 10). In group II, there was little bone
10 regeneration in the defect sites (Fig. 11). In Group III, hardly any bony regeneration was noted
11 (Fig. 12). The 3D CT image demonstrated some bony formation in Group IV at 6 weeks (Fig.
12 13A), and near-complete repair of the defective area at 3 months (Fig. 13B).

13 The H & E staining of Groups 1, 2 and 3 did not demonstrate good bone formation. In
14 Group IV, the experimental site showed cancellous bone formation (Fig. 14A), while the control
15 site did not (Fig. 14B). The Von Kossa stains of the experimental site were strongly positive for
16 mineralization, and no cortical formation at 3 months *in vivo* (Fig. 15).

17 The immunochemical stain for detecting adenovirus was positive in all pre-implantation
18 culture (Fig. 16A) and was negative in all 6 weeks and 3 months samples (Figs. 16B, 16C).

19 At 6 weeks, there were no statistically significant differences between the four groups.
20 At 3 months, in Group IV, there was a significant increase of bone formation of the defective site
21 induced by adenovirus bone morphogenetic protein (BMP)-2 transduction.

Table 1 shows the new bone formation areas by 3D CT (cm²). The statistical data revealed that Group IV, at 3 months, had significant increase of bone formation of the defective site induced by adenovirus bone morphogenetic protein (BMP)-2 transduction.

Table 1: New Bone Formation Areas by 3D CT (cm²)

Group (n=5)/area,cm2	Experimental	Control	
1 (6w)	3.31	2.13	0.3789
1 (3M)	4.51	2.87	0.2141
2 (6w)	3.17	2.35	0.2664
2 (3M)	4.17	3.97	ns
3 (6w)	1.37	1.01	ns
3 (3M)	4.17	2.72	ns
4 (6w)	4.32	3.61	ns
4 (3M)	6.32	4.41	p=0.0128

The constructed recombinant adenovirus with human BMP-2 gene successfully mediated the gene transfer and BMP-2 protein expression in the MSCs and produced an effective amount of the protein. Expressed BMP-2 protein was detected in culture medium of the transduced cell. The proper size of the detected protein suggested that adenovirus-mediated BMP-2 gene transfer and expression underwent the proper process in protein secretion and enzymolysis after translation. One band of BMP-2 protein has been observed in a Western blot analysis, and a similar pattern has also been found in a Chinese hamster ovary (CHO) cell (Cook, S.D., et al., 1994, *In Vivo Evaluation of Recombinant Human Osteogenic Protein Implants as a Bone Graft Substitute for Spinal Fusions*, SPINE 19:1655-1663) and osteosarcoma (Wang, E.A., et al., 1990, *Recombinant Human Bone Morphogenetic Protein Induces Bone Formation*, PROC NATL ACAD SCI U S A 87:2220-2224) expression media. This pattern may result from unclear proteinic enzymolysis.

1 Adenovirus-mediated BMP-2 gene transfer on MSCs directed the cells to proliferation
2 and differentiation. Cell response to the gene transfer may have been through an autocrine or
3 paracrine effect of BMP-2 protein that was expressed from the transferred BMP-2 gene by the
4 cell itself.

5 The results demonstrated that bone repair *in vivo* was induced and enhanced by
6 adenovirus bone morphogenetic protein (BMP)-2 transduction. Moreover, gene transfer has the
7 advantage of directing the cell to continuously produce BMP-2 protein. In an *in vitro* study, a
8 one-time transduction of recombinant adenovirus into the cells kept the cell exposure to BMP-2
9 until mineralization; this required continued incubation for as long as 30 days and several
10 medium changes (Lou, J., et al., 1999, *Gene Therapy: Adenovirus-Mediated Human Bone*
11 *Morphogenetic Protein-2 Gene Transfer Induces Mesenchymal Progenitor Cell Proliferation*
12 *and Differentiation In Vitro and Bone Formation In Vitro*, J. ORTHO. RES. 17:43-50). BMP-2
13 protein should continue to be released as long as the transduced gene stays in the cell.

14 Recombinant adenovirus-mediated gene transfer has its limitation in the immune-
15 competent host. The host immune response to the adenovirus proteins from leak expression of
16 virus genome will delete the recombinant adenovirus and terminate expression of the transgene.
17 Also, inflammation has been observed. Therefore, the use of a new generation of recombinant
18 adenovirus vectors that has less immunogenicity and longer expression in the immune-
19 competent host will help to reduce these problems and thus be more effective.

20 This study demonstrated that BMP-2 gene transfer into mesenchymal stem cells could
21 potentially direct differentiation into bone formation *in vivo*. Furthermore, this study
22 demonstrated that the use of the polymer Pancogene S/MSCs with *adv*-BMP-2 gene transfer will
23 enhance the bony healing of critical size craniofacial defect.

1 Table 2, below, shows the biomechanic test in Group IV experimental site compared with the
2 normal cranium bone.

3 **Table 2 Biomechanics test (MPa)**
4

normal cranial bone	88646.02 ±5121.803	P=0.3444
Group 4 (collagen)	80536.46 ±19302.64	

5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

1
2 **EXAMPLE 2:**

3 In a second example, the bone marrow stromal cells (MSC) were separated from the
4 aspirate from the iliac crest from 20 minipigs one month before implantation. The MSC were
5 expanded in monolayered culture. Full-thickness bone defects ($3 \times 1.2 \text{ cm}^2$) were created on the
6 bilateral maxilla of the minipigs. The osteogenic periosteum was removed. The experimental site
7 (on the animal's left), the MSC was implanted by using *ex vivo* adenovirus-BMP-2 (Bone
8 Morphogenic Protein-2) mediated gene transfer to the expanded MSC 5 days before implantation
9 with a concentration was $5 \times 10^7/\text{ml}$ of collagen type I (Pancogen). In the control site (on the
10 animal's right), the MSC was implanted by *ex vivo* adenovirus- β Gal medicated gene transfer
11 under the same condition. After 3 months, 20 minipigs were sacrificed, the head sent for 3D CT
12 examination and then sawed in half. One half was preserved in -80°C refrigerator for further
13 biomechanic testing. The other half was sawed into slices for histology (HE and Von Kossa
14 stain) and immunochemical stain for detection of adenovirus.

15 Near-complete maxilla defect repair was achieved by simple aspiration of bone marrow
16 stromal cells and after three months implantation of tissue engineered constructs with collagen
17 type I. The histology showed mature woven bone formation. The regenerated bone and the
18 biomechanic test are similar to that of normal cranial bone. This is a cell therapy which is
19 further based on *ex vivo* infection, thus the gene is securely delivered to target cells and the viral
20 contact to the host animal will be significantly reduced.

MATERIALS AND METHODS

Construction of Recombinant Adenovirus

Recombinant adenovirus BMP-2 was constructed by *in vivo* homologous recombination in 293 cells between expression vector pAC-cytomegalovirus BMP-2 and the large Cla I fragment (2.6-100 mm) of adenovirus type 5 genome DNA. Briefly, to make recombinant plasmid pAC-cytomegalovirus BMP-2, the human BMP-2 gene (no. 40345; ATCC [American Type Culture Collection], Rockville, MD, U.S.A.) was inserted into the vector pAC cytomegalovirus. The resulting plasmid was made linear by Xho I and cotransfected with the large Cla I fragment of wild-type adenovirus type 5 genome DNA by lipofectamine (GIBCO BRL, Grand Island, NY, U.S.A.). For each 35-mm dish of 293 cells, 2 mg of each kind of DNA (total: 4 mg) was transfected. Plaques containing recombinant adenovirus were picked after 10 days of agarose overlay and incubation at 37°C. The viral stocks were propagated and titered in 293 cells. The recombinant adenovirus with human BMP-2 gene, which was under transcriptional control of the cytomegalovirus early gene promoter/enhancer, was named adenovirus BMP-2. The recombinant adenovirus containing the bacterial β -galactosidase (*lac z*) gene was constructed by the same procedure, and the virus was named adenovirus β -galactosidase. Viral stock of 293 lysate was titered in 293 cells by serial dilution and plaque-formation assay.

Western Blot Detection of Adenovirus BMP-2 Expression in MSC (Bone Marrow Stromal Cells) *In Vitro*

Cells from MSC (Bone Marrow Stromal Cells) were grown in Dulbecco's modified Eagle medium with 10% fetal calf serum to 90% confluence. The cells were infected with adenovirus β -galactosidase or adenovirus BMP-2 at a multiplicity of infection of 50. The infected cells were maintained in Dulbecco's modified Eagle medium with 10% fetal calf serum

1 and 50 mg/ml ascorbic acid. The medium was collected 7 days later and then
2 immunoprecipitated with monoclonal anti-body h4b2/5.10.24 (a gift from Genetics Institute,
3 Cambridge, MA, U.S.A.). The immunoprecipitation was then run on a 12.5% sodium dodecyl
4 phosphate-polyacrylamide gel electrophoresis and transferred overnight to a polyvinylidene
5 difluoride membrane (Millipore, Bedford, MA, U.S.A.). The expressed protein was detected with
6 monoclonal antibody h4b2/5.10.24 and goat anti-mouse IgG alkaline-phosphatase conjugate
7 (Sigma Chemical, St. Louis, MO, U.S.A.).

8 **Bone Marrow Aspiration and Adenoviral Infection**

9 20 female minipigs (Mitsae Pig, CGU, Taiwan) were used in this experiment. The
10 minipigs were sacrificed at 3 months after tissue engineered constructs implantation. At the
11 first stage of surgery, bone marrow aspiration was performed when minipigs were under
12 intramuscular injection of 1 ml/ 10 kg 2% Rompum (Bayer, USA), Ketamine injection
13 (50mg/ml, Yung Shin Pharmaceutical Industrial Co., Taiwan). 20 cc of the aspirated bone
14 marrow was mixed with 2 cc of heparin Sodium (5000 unit/ml, Agglutex, China Chemical &
15 Pharmaceutical Co., Taiwan). Aspirated iliac crest marrow (5-10 ml) was transferred to sterile
16 tubes, to which 20 ml complete medium was added. The tubes were spun at 1000 rpm for 5 min
17 to pellet the cells, the supernatants and fat layers were removed, and the cell pellets (2.5-5.0 ml)
18 were resuspended and loaded onto 70% Percoll (Sigma, St. Louis, Missouri) gradients. These
19 gradients were centrifuged at 460 x g for 15 min and harvested into three fractions with a pipet:
20 the top 25% of the gradients (low density cells), pooled density = 1.03 g per ml; the middle 50%
21 of the gradients (high density cells), pooled density = 1.10 g per ml; density = 1.14 g per ml. In
22 preliminary experiments, each of these three pools was plated separately in complete medium in
23 100 mm dishes. Adherent marrow-derived mesenchymal cells were observed to be localized to
24 the low-density fraction. To produce adherent cell cultures for all subsequent experiments, only

1 the cells in the low density fraction were plated. The medium was changed every 3-4 days. After
2 one month, an adequate number of cells was reached (Fig. 1).

3 At the experimental site, Adenovirus mediated human BMP-2 gene transfer to MSCs was
4 performed 5 days before the secondary stage of surgery (Fig. 1). As a control, Adenovirus
5 mediated β Gal gene transfer was carried out at the same time and for the same period of time.
6 The cells were trypsinized by 25% trypsin and washed for three times. Cell numbers were
7 determined with a hemocytometer. The cells were then mixed with four different polymers to
8 achieve the same concentration of $50 \times 10^6/\text{ml}$. After one hour in the incubator, the cells with the
9 polymer Pancogene S became more gel-like.

10 **Open Implantation on the Maxilla Defects After Creating Maxillary Bone Defect**

11 At the secondary stage of surgery, all minipigs were intubated and on ventilators.
12 Bilateral maxillary defects of $3 \times 1.2 \text{ cm}^2$ each were created by removal of bone and periosteum,
13 confirming that no osteoinductive tissue exists (Fig. 2 A,B). To bridge the defect, one
14 biodegradable PLLA 6-holes miniplate was used on each maxillary defect (Fig. 2C). Each 3ml
15 of cell/polymer construct was implanted to each site (Fig. 2D). The right sided defect was
16 covered with adenovirus-mediated BMP-2 gene transferred MSCs/polymer. The left side defect
17 was filled with adenovirus-mediated β Gal gene transferred on MSCs/polymer. The facial wounds
18 were closed in a watertight manner.

19 **Histologic Examination of Bone Samples Harvested from the Maxilla Defects**

20
21
22 Harvested samples were fixed in buffered 10% formalin for 72 hours and then sawed in
23 half. Half of them were decalcified in Decalcifier I (Surgipath, Northbrook, IL, U.S.A.) solution
24 for another 48 hours. The samples then underwent procedures for embedding, incising, and

1 staining with hematoxylin and eosin. The nondecalcified other half of the samples was stained
2 with von Kossa's silver nitrate for demonstration of matrix mineralization.

3 **Immunohistochemistry for Adenovirus Hexon Protein**

4 Immunohistochemistry was performed using formalin-fixed, paraffin embedded tumor
5 sections. After removal from paraffin, slides were hydrated and digested with pronase (Meridian
6 Diagnostics, Cincinnati, OH). The primary antibody (MAB805, Chemicon International,
7 Temecule, CA) was applied at 1:500 dilution, and slides were incubated for 1 hr at 35 °C. A
8 biotinylated goat anti-rabbit secondary was then applied, followed by a streptavidin-horseradish
9 peroxidase conjugate. Diaminobenzidine (DAB) was used as the chromogen, before slides were
10 counterstained with hematoxylin.

11 **Biomechanical Analysis**

12 The samples were stored at -70°C before testing. All the samples of engineered bones
13 were sawed using a 6-mm dermal punch to form disks. Each disk was immediately mounted in
14 an electrically insulating cylindrical confining chamber. The chamber was mounted in a servo-
15 controlled Dynastat mechanical spectrometer (IMASS, Hingham, MA) interfaced to a computer.
16 The samples were equilibrated at room temperature in 0.15M PBS, pH 7.4, containing 100
17 units/ml penicillin G and 100 ug/ml streptomycin. Samples were compressed between a porous
18 polyethylene platen and at the base of the chamber.

19 After mounting each disk in the confined compression chamber, the distance between the
20 porous platen and the chamber was decreased until a signal of ~5 gm was detected by the load
21 cell. This distance was taken to be the sample thickness. Each disk was compressed by 10
22 sequential increments of 2.5-3.0% static strain, up to a maximum of 25-30% total strain. After
23 each increment, the load was recorded every 0.5 seconds for 100 seconds. Stress relaxation data

1 was fit to a poroelastic model of material behavior that yielded values for the material properties;
2 equilibrium modulus.

3 **Statistical Analysis**

4 All measurements were collected in triplicate and expressed as means \pm standard
5 deviations. Single factor analysis of variance (ANOVA) was employed to assess the statistical
6 significance of results for all the 3D CT images of the repaired bone areas.

7 **RESULTS**

8 **Recombinant Adenovirus Construction and Human BMP-2 Expression**

9 Adenovirus BMP-2, a replication-defective adenovirus vector encoding human BMP-2,
10 was constructed. In this vector, transcription of the BMP gene is controlled by cytomegalovirus
11 early gene promoter/enhancer. The recombinant adenovirus encoding *Escherichia coli* lac z gene
12 (adenovirus β -galactosidase) was constructed by the same procedure as the control vector. These
13 two recombinant adenoviruses have been deleted only in the E1 region of the genome. The E3
14 region of the virus genome was purposefully retained intact on the vectors because of evidence
15 that the E3 proteins of the virus inhibit the host immune response. Titered in 293 cells, the viral
16 lysate of the vectors are usually 10^8 plaque-forming units per milliliter.

17 The gene expression of human BMP-2 protein in MSCs, which was mediated by
18 adenovirus BMP-2 gene transfer, was assessed by detecting the coding product with monoclonal
19 antibody h4b2/5.10.24 in transduced cells. The secreted active BMP-2 molecules were detected
20 in cultured medium from adenovirus BMP-2-transduced MSCs by immunoprecipitation and
21 immunoblot analysis but were not found in cultured medium from either adenovirus β -
22 galactosidase-infected MSCs. The BMP gene transfected by adenovirus seems well expressed
23 and processed in MSCs. The molecular weight of secreted BMP-2 indicated mature status

1 without prepeptide. The band was noticeable at 18KDa due to overlap of bands at 16 and 22
2 KDa.

3 There was noticeable whitish bone formation at the site infected with BMP-2 at
4 3 months, while no obvious bone formation was found at the β Gal infection site (Fig. 3).

5 3D-CT image studies were performed for each sample. The 3D CT image demonstrated
6 complete repair of the defect area at 3 month at the experimental site (Fig. 4).

7 The H & E staining of the experimental site showed cancellous bone formation, while the
8 control site did not (Fig. 5). The Von Kossa stain of the experimental site were strongly positive
9 for mineralization, and mature woven *de novo* bone formation at 3 months *in vivo* (Fig. 5).

10 At 3 months, there was a significant increase of bone formation of the defect site induced
11 by adenovirus bone morphogenetic protein (BMP)-2 transduction as shown in Table 3, below.

12 **Table 3- New Bone Formation Areas by 3D CT (n =10) (cm²)**

	Mean +/- SD
Experimental	2.6275 +/- 0.1795
Control	1.78 +/- 0.0764

13 p-value = 0.0154

14 The biomechanical properties between the harvest in the experimental site and normal
15 maxilla bone did not show statistical difference.

16 The contents of all cited references throughout this application are hereby expressly
17 incorporated by reference. The practice of the present invention will employ, unless otherwisc
18 indicated, conventional techniques of pharmacology and pharmaceuticals, which are within the
19 skill of the art.

6

Parameter	Unit	Value	Uncertainty
α	deg	10.0	0.5
β	deg	10.0	0.5
γ	deg	10.0	0.5
δ	deg	10.0	0.5
ϵ	deg	10.0	0.5
ζ	deg	10.0	0.5
η	deg	10.0	0.5
θ	deg	10.0	0.5
ϕ	deg	10.0	0.5
χ	deg	10.0	0.5
ψ	deg	10.0	0.5
ω	deg	10.0	0.5
ν	deg	10.0	0.5
μ	deg	10.0	0.5
λ	deg	10.0	0.5
κ	deg	10.0	0.5
ι	deg	10.0	0.5
\hbar	deg	10.0	0.5
g	deg	10.0	0.5
f	deg	10.0	0.5
e	deg	10.0	0.5
d	deg	10.0	0.5
c	deg	10.0	0.5
b	deg	10.0	0.5
a	deg	10.0	0.5